

NMSA405: exercise 8 – optional sampling theorem

Exercise 8.1: (Proposition 3.5) Under the assumptions of the Theorem 3.4 show that the condition

$$X_T^+ \in L_1 \quad \text{and} \quad \int_{[T > n]} X_n^+ d\mathbb{P} \xrightarrow{n \rightarrow \infty} 0$$

is equivalent to the uniform integrability of the sequence $(X_{T \wedge n}^+)$. Recall that $T < \infty$ almost surely.

Exercise 8.2: Let (X_n) be a sequence of iid random variables with $\mathbb{P}(X_1 = 1) = \mathbb{P}(X_1 = -1) = 1/2$ and let $S_n = \sum_{k=1}^n 2^{k-1} X_k$, $n \in \mathbb{N}$. Consider the first hitting time T of the sequence (S_n) of the set $\{1\}$. Then for (S_n) and T the optional sampling theorem does not hold. Show that $\mathbb{E}S_1 \neq \mathbb{E}S_T$ and the condition $\lim_{n \rightarrow \infty} \int_{[T > n]} |S_n| d\mathbb{P} = 0$ is not fulfilled.

Exercise 8.3: (remark to the Theorem 3.6) Let (X_n) be a \mathcal{F}_n -martingale and $T < \infty$ a.s. be a \mathcal{F}_n -stopping time. Show that the condition

$$\exists 0 < c < \infty : T > n \implies |X_n| \leq c \quad \text{a.s.}$$

does not imply the condition

$$X_T \in L_1 \quad \text{and} \quad \int_{[T > n]} |X_n| d\mathbb{P} \xrightarrow{n \rightarrow \infty} 0$$

from the Theorem 3.4.

Hint: Consider the sequence $X_n = \sum_{k=1}^n 3^k Y_k$ where (Y_k) is a sequence of iid random variables with the uniform distribution on $\{-1, 0, 1\}$.

NMSA405: exercise 9 – discrete and non-trivial random walk

Definition: Let (X_n) be an iid random sequence such that $\mathbb{P}(X_1 = 1) = p$ and $\mathbb{P}(X_1 = -1) = 1 - p$ where $p \in [0, 1]$. We call the corresponding random walk (S_n) a (*simple*) *discrete random walk*. If $p = 1/2$ we get the symmetric simple random walk.

Exercise 9.1: Consider the stopping time $T^B = \min\{n \in \mathbb{N} : S_n \notin B\}$ defined as the first exit time of the discrete random walk S_n from the bounded set $B \in \mathcal{B}(\mathbb{R})$ and the stopping time $T_a = \min\{n \in \mathbb{N} : S_n = a\}$ defined as the first hitting time of the random walk S_n of the set $\{a\}$ for $a \in \mathbb{Z}$. Show that

1. $T^B < \infty$ a.s.,
2. $T_a < \infty$ a.s. if $p = 1/2$.

Exercise 9.2: Show that the discrete random walk fulfills

- (i) $S_n \xrightarrow{n \rightarrow \infty} \infty$ a.s. $\iff p > 1/2$,
- (ii) $S_n \xrightarrow{n \rightarrow \infty} -\infty$ a.s. $\iff p < 1/2$,
- (iii) $\limsup_{n \rightarrow \infty} S_n = \infty$ a.s., $\liminf_{n \rightarrow \infty} S_n = -\infty$ a.s. $\iff p = 1/2$.

Exercise 9.3: Consider a discrete symmetric random walk (S_n) . For $a, b \in \mathbb{Z}$, $a < 0$, $b > 0$, we define $T_{a,b} = \min\{n \in \mathbb{N} : S_n \notin (a, b)\}$ as the first exit time of S_n from the interval (a, b) . Show that in that case

$$\mathbb{P}(S_{T_{a,b}} = a) = \frac{b}{b-a} \quad \text{and} \quad \mathbb{E}T_{a,b} = -ab.$$

Corollary: (i) $\mathbb{E}T^B < \infty$ for any bounded set $B \in \mathcal{B}(\mathbb{R})$, (ii) $\mathbb{E}T_b = \infty$ for any $b \in \mathbb{Z}$, $b \neq 0$.

Exercise 9.4: Let (S_n) be a symmetric simple random walk and let $A < 0 < B$ be independent integrable random variables, independent of (S_n) . Denote $T = \min\{n \in \mathbb{N} : S_n \notin (A, B)\}$. Show that in that case

$$\mathbb{P}(S_T = A) = \mathbb{E} \frac{B}{B-A} \quad \text{and} \quad \mathbb{E}T = -\mathbb{E}A \cdot \mathbb{E}B < \infty.$$

Definition: The random sequence $S_n = \sum_{k=1}^n X_k$, $n \in \mathbb{N}$, where (X_n) is an iid random sequence and $\mathbb{P}(X_1 \neq 0) > 0$ is called a *non-trivial random walk*.

Theorem 3.11: Let T^B be the first exit time of the non-trivial random walk (S_n) from the bounded set $B \in \mathcal{B}(\mathbb{R})$. Then $\mathbb{E}(T^B)^r < \infty$ for any $r \in \mathbb{N}$.

Idea of proof: There is $m \in \mathbb{N}$ such that $p = \mathbb{P}(|S_m| > d) > 0$ where d is the diameter of the set $B \cup \{0\}$. Consider the stopping time $\tau = \inf\{k \in \mathbb{N} : |S_{mk} - S_{m(k-1)}| > d\}$. Then τ has a shifted geometrical distribution with parameter p and $T^B \leq m\tau$.

NMSA405: exercise 10 – convergence theorems

Exercise 10.1: Give an example of a martingale which converges to the random variable $X_\infty \in L_1$ almost surely but not in L_1 .

Exercise 10.2: Let (Y_n) be a sequence of independent random variables such that

$$\mathbb{P}(Y_n = 2^n - 1) = 2^{-n}, \quad \mathbb{P}(Y_n = -1) = 1 - 2^{-n}, \quad n \in \mathbb{N}.$$

Check that $X_n = \sum_{k=1}^n Y_k$ is a martingale. Show that $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} -\infty$ and hence the assumptions of the martingale convergence theorems cannot be fulfilled.

Exercise 12.3: (martingale proof of the Kolmogorov 0-1 law) Let $X = (X_1, X_2, \dots)$ be a sequence of independent random variables and $F = [X \in T]$ where $T \in \mathcal{T}$ is a terminal set. Show that

$$\forall n \in \mathbb{N} \quad \mathbb{E}[\mathbf{1}_F | \mathcal{F}_n] = \mathbb{P}(F) \quad \text{a.s.} \quad \text{and at the same time} \quad \mathbb{E}[\mathbf{1}_F | \mathcal{F}_n] \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \mathbf{1}_F.$$

From this conclude that $\mathbb{P}(F)$ is either 0 or 1.