

Proof of Proposition 1.19

$X \text{ - HLCs, } \dim X = n$

$K \subset X$ compact convex $\Rightarrow \forall x \in K : x \text{ is a convex combination}$
of points $x_1, \dots, x_k \in X$ with properties:

- $x_1, \dots, x_k \in \text{ext } K$
- x_1, \dots, x_k are affinely independent
- $k \leq n+1$

Recall: x_1, \dots, x_k are affinely independent $\Leftrightarrow x_2 - x_1, \dots, x_k - x_1$ are
linearly independent.

Proof ① $n=0 \Rightarrow k=\{0\}$

$n=1 \Rightarrow k=\{0, 1\} \subset \mathbb{R}$ or $k=\{1\} \subset \mathbb{R}$

clear

② Suppose $n \geq 1$ and that it is true if $\dim X \leq n$.

Suppose $\dim X = n+1$

- WLOG $0 \in K$ (replace K by $K+x$ for some $x \in K$)
- WLOG $\text{span } K = X$ (otherwise replace X by $\text{span } K$
and apply the induction hypothesis
for $\dim(\text{span } K) \leq n$)
- Then K has nonempty interior

For $x_1, \dots, x_{n+1} \in K$ be linearly independent,
have a basis of X

The $T: \mathbb{R}^{n+1} \rightarrow X$ defined by

$T(t_1, \dots, t_{n+1}) = \sum_{j=1}^{n+1} t_j \cdot x_j$ is an isomorphism
of \mathbb{R}^{n+1} onto X

Then we deduce that $\text{co}\{0, x_1, \dots, x_{n+1}\}$ has
nonempty interior, as on \mathbb{R}^{n+1}
 $\text{co}\{0, e^1, \dots, e^{n+1}\}$ has nonempty interior

So, $\text{int } K \neq \emptyset$

• Let $x \in K$. If $x \in \partial K$, then $x \notin \text{bulk } K$,
so $\exists f \in \mathcal{F}^*$ $f(x) \geq \sup_{\text{bulk } K} f = \sup_{K} f$

So, $F = \{y \in K, f(y) = f(x)\}$
is a closed face

bulk K is dense in K
(see Lemma V.16)

We can use induction hypothesis
on F (its dimension is at most n) [and L17(a, s)]

* $x \in \text{bulk } K$ • Fix $y \in \text{ext } K$ arbitrary. Consider the
line $y + t(x-y)$, $t \in \mathbb{R}$. This line intersects K in a segment
one endpoint is y . Denote the second one by z . Then
 $z \in \partial K$. Use the previous case to deduce that

$z = t_1 a_1 + \dots + t_k a_k$, $a_1, \dots, a_k \in \text{ext } K$ (in fact,
they are in $\text{ext } F$, where F is the face constructed
in the previous case), $k \leq n$, a_1, \dots, a_k
affinely independent.

Since $x = (1-s)y + s z$ for some $s \in (0, 1)$, we have
 $x = (1-s)y + s t_1 a_1 + \dots + s t_k a_k$

It remains to observe that y, a_1, \dots, a_k are affinely independent.

Let $t_0(y-a_1) + t_1(a_2-a_1) + \dots + t_k(a_k-a_1) = 0$

Applying the function f constructed in the first case so

that $f(a_k) = f(a_2) = \dots = f(a_k) > f(y)$

Then $t_0(f(y) - f(a_1)) = 0$, so $t_0 = 0$

Since a_1, \dots, a_k are aff. independent, we see

$$a_2 = \dots = a_k = 0$$